



From Seed to Harvest: How Plant Hormones Drive Growth and Agricultural Innovation

Achanta Rama*

Principal, BVK College, Visakhapatnam (Andhra Pradesh), India.

(Corresponding author: Achanta Rama)*

(Received: 03 Feb. 2025; Revised: 16 April 2025; Accepted: 03 May 2025; Published: 20 May 2025)

(Published by Research Trend)

DOI: <https://doi.org/10.65041/abi.16>

ABSTRACT: Plant hormones, or phytohormones, are essential chemical messengers that regulate the intricate processes of plant growth, development, and environmental adaptation. These signaling molecules, including auxins, gibberellins, cytokinins, abscisic acid (ABA), and ethylene, orchestrate critical stages from seed germination to fruit ripening, ensuring plants thrive under diverse conditions. This research article explores the multifaceted roles of plant hormones, their interactions, and their transformative applications in modern agriculture, while addressing challenges and future directions for sustainable innovation. Auxins drive cell elongation, root development, and apical dominance, with synthetic analogs like 2, 4-dichlorophenoxyacetic acid (2, 4-D) enhancing tissue culture and rooting in crop propagation. Gibberellins promote stem elongation, seed germination, and flowering, widely used to increase fruit size in crops like grapes and citrus and to break seed dormancy in cereals. Cytokinins stimulate cell division, organogenesis, and delay senescence, improving crop yields and extending post-harvest shelf life in leafy greens and tomatoes. ABA governs stress responses, seed dormancy, and stomatal regulation, with applications in developing drought-resistant crops and mitigating abiotic stresses. Ethylene, a gaseous hormone, regulates fruit ripening, leaf abscission, and stress adaptation, with ethylene-releasing compounds like ethephon synchronizing ripening and inhibitors like 1-methylcyclopropene (1-MCP) prolonging storage life. Hormonal crosstalk, such as the interplay between auxins and cytokinins in shoot-root balance or gibberellins and ABA in germination, ensures precise developmental control, offering opportunities for targeted agricultural interventions. In agriculture, synthetic hormones and biotechnological advances, including CRISPR-Cas9 gene editing, have revolutionized crop improvement by enhancing yields, stress tolerance, and fruit quality. For instance, ABA-based treatments improve drought resistance in wheat and maize, while gibberellin and cytokinin applications boost fruit set in horticultural crops. However, challenges persist, including the environmental risks of synthetic hormone overuse, such as herbicide runoff and soil microbial disruption, and the need for precise application to avoid growth abnormalities. Future research should prioritize precision hormone engineering, leveraging synthetic biology, nanotechnology, and digital agriculture to develop eco-friendly hormone analogs and optimize delivery systems. These innovations promise to minimize off-target effects and enhance sustainability. By integrating hormone regulation with climate-resilient crop varieties, agriculture can address global food security amidst challenges like climate change and population growth. This article underscores the pivotal role of plant hormones in driving agricultural productivity and resilience, highlighting the need for continued research to unlock their full potential. Sustainable hormone-based strategies will be crucial for transforming agriculture, ensuring robust crop performance from seed to harvest while safeguarding environmental health for future generations.

Keywords: Plant hormones, agriculture, growth, innovation, sustainability.

INTRODUCTION

Plant hormones, or phytohormones, are chemical messengers that orchestrate the growth,

development, and environmental responses of plants. These signaling molecules regulate critical processes such as cell division, elongation, differentiation, and stress adaptation, enabling

plants to thrive in diverse conditions (Davies, 2010). From seed germination to fruit ripening, phytohormones ensure plants complete their life cycles while responding to external cues like light, temperature, and water availability. In agriculture, understanding and manipulating plant hormones have revolutionized crop production, enhancing yields, improving stress tolerance, and extending post-harvest shelf life (Santner & Estelle 2009). The significance of plant hormones extends beyond natural growth processes to their transformative role in modern agriculture. By leveraging phytohormones, farmers and scientists can optimize crop traits, mitigate environmental stresses, and address global challenges like food security and climate change (Peleg & Blumwald 2011). This paper aims to explore the major classes of plant hormones, their functions, their interactions, and their applications in agriculture. It also discusses challenges and future directions for hormone-based innovations, emphasizing sustainable agricultural practices.

LITERATURE REVIEW

Plant hormones, or phytohormones, are pivotal in regulating plant growth and development, influencing processes such as seed germination, root and shoot development, flowering, and fruit ripening (Thapa *et al.*, 2024). Key hormones include auxins, gibberellins, cytokines, abscisic acid, and ethylene, each playing distinct roles; for instance, auxins promote cell elongation while cytokinins stimulate cell division (Csukasi *et al.*, 2009). The intricate interactions among these hormones form complex networks that enable plants to adapt to environmental changes and stressors (Choudhary & Kumari 2021). Advances in understanding hormone biosynthesis and signaling pathways have led to biotechnological innovations, allowing for the development of transgenic crops with enhanced traits, thereby improving agricultural productivity and sustainability (Davies, 1987). This knowledge not only enhances basic plant science but also offers practical applications in crop management, ultimately addressing global challenges such as food security and climate change (Nagel, 2024; Csukasi *et al.*, 2009). Indirect mechanisms involve the suppression of phytopathogens through the production of antimicrobial compounds, competition for resources and induction of systemic resistance in plants (Rathod *et al.*, 2025). Maximum leaf area, fruit weight and minimum days to flowering were observed under the foliar application of NAA 50 ppm. Overall, it was concluded that GA3 75 ppm was best in improving shoot, root and yield attributes of strawberry (Singh *et al.*, 2023).

MAJOR CLASSES OF PLANT HORMONES AND THEIR FUNCTIONS

A. Auxins

Auxins, primarily indole-3-acetic acid (IAA), are pivotal in regulating plant growth. They promote cell elongation, root initiation, and apical dominance, where the main shoot suppresses lateral bud growth (Gray, 2004). Auxins influence tropisms, such as phototropism and gravitropism, guiding plants toward light or gravity. In agriculture, synthetic auxins like 2, 4-dichlorophenoxyacetic acid (2, 4-D) are used in tissue culture to induce callus formation and in rooting powders to stimulate adventitious root growth in cuttings (Davies, 2010). These applications enhance propagation efficiency for crops like grapes and ornamentals.

B. Gibberellins

Gibberellins (GAs) regulate stem elongation, seed germination, and flowering. They break seed dormancy by stimulating enzyme production, such as amylase, which mobilizes stored nutrients (Hedden & Thomas 2012). In agriculture, gibberellins are applied to increase fruit size in seedless grapes and citrus, promote uniform flowering in cereals, and enhance malting in barley. Their ability to elongate stems is exploited to improve plant architecture, making crops like rice more resistant to lodging (Sasaki *et al.*, 2002). GAs have been linked to improved crop yields and quality in horticultural plants, enhancing traits like flowering time and fruit development, which is crucial for agricultural productivity (Lupepsa *et al.*, 2024).

C. Cytokinins

Cytokinins (CKs), such as zeatin, promote cell division, organogenesis, and delay senescence (Kieber & Schaller 2014). They work synergistically with auxins to regulate tissue differentiation in vitro, enabling the regeneration of whole plants from explants. In agriculture, cytokinins enhance crop yields by stimulating shoot growth and increasing grain filling in cereals. They also extend post-harvest shelf life by delaying leaf and fruit senescence, benefiting crops like leafy greens and tomatoes. CKs exhibit spatial distribution within plant tissues, with transport facilitated by specific transporters, which is essential for their function in response to environmental cues (Zhao *et al.*, 2024).

D. Abscisic Acid (ABA)

Abscisic acid (ABA) is a stress-response hormone that regulates seed dormancy, stomatal closure, and abiotic stress tolerance (Cutler *et al.*, 2010). ABA is transported from the shoot to the fruit, where it plays a significant role in maturation and ripening, often interacting synergistically with ethylene to regulate these processes (Gupta *et al.*, 2022). Understanding ABA's diverse roles and mechanisms can lead to applications in agriculture and medicine, enhancing crop yield and quality and offering therapeutic benefits in human health.

(Lievens *et al.*, 2017; Gupta *et al.*, 2022). During drought or salinity stress, ABA induces stomatal closure to reduce water loss and activates stress-responsive genes. In agriculture, ABA analogs are used to develop drought-resistant crops, such as wheat and maize, and to prolong seed dormancy in storage, preventing pre-harvest sprouting. ABA's role in stress mitigation is critical for sustaining yields in water-scarce regions (Zhu, 2016).

E. Ethylene

Ethylene, a gaseous hormone, regulates fruit ripening, leaf abscission, and stress responses (Bleecker & Kende 2000). It triggers the ripening of climacteric fruits like bananas and tomatoes by upregulating genes for softening and sugar accumulation. Ethylene also mediates responses to flooding and pathogen attack. In agriculture, ethylene-releasing compounds like ethephon are used to synchronize fruit ripening and facilitate mechanical harvesting. Conversely, ethylene inhibitors, such as 1-methylcyclopropene (1-MCP), extend storage life by delaying ripening.

HORMONAL CROSSTALK AND REGULATION

Plant hormones do not function in isolation; their interactions, or crosstalk, fine-tune growth and development (Santner & Estelle 2009). Auxins and cytokinins exhibit antagonistic effects in shoot and root development. High auxin-to-cytokinin ratios favor root formation, while high cytokinin-to-auxin ratios promote shoot growth (Kieber & Schaller 2014). This balance is exploited in tissue culture to regenerate specific organs. Gibberellins and ABA interact in seed germination, where gibberellins promote embryo growth, and ABA maintains dormancy (Hedden & Thomas 2012). Environmental cues, such as light or moisture, shift this balance to initiate germination.

Ethylene and auxin crosstalk regulates senescence and abscission. Auxin inhibits ethylene production to delay leaf drop, while ethylene accelerates abscission in aging tissues (Bleecker & Kende 2000). These interactions are mediated by complex signaling pathways involving transcription factors and secondary messengers. Understanding hormonal crosstalk is essential for developing targeted agricultural interventions, as manipulating one hormone often affects others, impacting overall plant performance.

AGRICULTURAL APPLICATIONS OF PLANT HORMONES

A. Synthetic Plant Hormones in Crop Improvement
Synthetic hormones have transformed agriculture by enabling precise control over plant growth. Synthetic auxins, such as 2,4-D, are used as herbicides to selectively eliminate broadleaf weeds while sparing grasses (Davies, 2010). Gibberellin applications increase fruit size and uniformity, benefiting horticultural crops like apples and pears

(Hedden & Thomas 2012). Cytokinin-based treatments enhance grain yields in rice and wheat by promoting sink strength, ensuring efficient nutrient allocation to developing seeds (Kieber & Schaller 2014).

B. Stress Resistance

Phytohormones play a critical role in enhancing stress tolerance. ABA applications improve drought and salinity tolerance by upregulating stress-responsive genes and reducing transpiration (Cutler *et al.*, 2010). Transgenic crops overexpressing ABA biosynthesis genes exhibit enhanced survival under water-limited conditions (Zhu, 2016). Ethylene modulates responses to biotic stresses, such as pathogen infections, by inducing defense-related proteins (Bleecker & Kende 2000). These advancements are vital for sustaining agriculture in marginal lands affected by climate change.

C. Enhancing Fruit Yield and Quality

Hormonal manipulation improves fruit yield and quality. Gibberellins and cytokinins increase fruit set and size in crops like strawberries and kiwifruit (Hedden & Thomas 2012; Kieber & Schaller 2014). Ethylene management ensures uniform ripening, critical for supply chain efficiency (Bleecker & Kende 2000). Post-harvest treatments with cytokinins or ethylene inhibitors maintain fruit firmness and flavor, reducing spoilage and extending marketability. These strategies are particularly valuable for perishable crops in developing nations with limited cold storage.

D. Biotechnological Innovations

Biotechnology has unlocked new possibilities for hormone regulation. Genetic engineering of hormone biosynthesis or signaling pathways creates crops with desired traits, such as dwarfism for lodging resistance or enhanced stress tolerance (Sasaki *et al.*, 2002). CRISPR-Cas9 enables precise editing of hormone-related genes, offering a more targeted approach than traditional breeding. For example, modifying gibberellin oxidase genes produces semi-dwarf rice varieties with higher yields. These innovations promise to address global food demands sustainably (Peleg & Blumwald 2011).

CHALLENGES AND FUTURE DIRECTIONS

A. Limitations of Hormone-Based Applications

Despite their benefits, hormone-based applications face challenges. The efficacy of synthetic hormones depends on environmental conditions, application timing, and crop species, requiring precise management (Davies, 2010). Over application can disrupt hormonal balance, leading to abnormal growth, reduced yields, or environmental contamination. For instance, excessive auxin use in tissue culture can inhibit shoot regeneration, while gibberellin overuse may

cause overly elongated, weak stems (Hedden & Thomas 2012).

B. Risks of Synthetic Hormone Overuse

The widespread use of synthetic hormones raises ecological concerns. Runoff from auxin-based herbicides can contaminate water bodies, harming non-target plants and aquatic ecosystems (Davies, 2010). Prolonged exposure to synthetic hormones may also induce resistance in weeds or alter soil microbial communities. Regulatory frameworks must balance agricultural benefits with environmental safety, emphasizing biodegradable and eco-friendly hormone analogs (Zhu, 2016).

C. Future Research on Precision Hormone Engineering

Future research should focus on precision hormone engineering to enhance sustainability. Advances in synthetic biology can develop hormone analogs with targeted effects, minimizing off-target impacts (Peleg & Blumwald 2011). Nanotechnology offers potential for controlled hormone delivery, ensuring optimal concentrations at specific plant tissues. Integrating hormone regulation with digital agriculture, such as sensor-based monitoring of plant stress, can optimize application timing and dosage. Additionally, exploring natural hormone mimics from microbes or algae could reduce reliance on synthetic compounds (Santner & Estelle 2009).

CONCLUSIONS

Plant hormones are indispensable drivers of growth, development, and environmental adaptation, shaping the trajectory from seed to harvest. Auxins, gibberellins, cytokinins, ABA, and ethylene collectively regulate critical processes, with their crosstalk ensuring precise control (Santner & Estelle 2009). In agriculture, phytohormones enhance crop yields, stress tolerance, and post-harvest quality, addressing global food security challenges (Peleg & Blumwald 2011). Synthetic hormones and biotechnological innovations have expanded these applications, but challenges like overuse and environmental risks necessitate sustainable solutions. Continued research into precision hormone engineering, eco-friendly analogs, and integrated agricultural technologies will unlock new frontiers. By harnessing the power of plant hormones, we can develop resilient, high-yielding crops to feed a growing population while safeguarding the environment. The future of agriculture lies in understanding and manipulating these chemical messengers, ensuring a sustainable path from seed to harvest.

REFERENCES

- Bleecker, A. B. & Kende, H. (2000). Ethylene: A gaseous signal molecule in plants. *Annual Review of Cell and Developmental Biology*, 16, 1-18.
- Cutler, S. R., Rodriguez, P. L., Finkelstein, R. R. & Abrams, S. R. (2010). Absciscic acid: Emergence of a core signaling pathway. *Annual Review of Plant Biology*, 61, 651-679.
- Choudhary, M. & Kumari, A. (2021). Understanding Plant Hormones: Mechanisms and Functions in Growth and Development. *Plant Science Archives*, 6(1), 14-16.
- Csukasi, F., Merchante, C. & Valpuesta, V. (2009). Modification of plant hormone levels and signaling as a tool in plant biotechnology. *Biotechnology Journal*, 4(9), 1293-1304.
- Davies, P. J. (1987). Plant hormones and their role in plant growth and development. Springer.
- Davies, P. J. (2010). *Plant Hormones: Biosynthesis, Signal Transduction, Action!* Springer.
- Gray, W. M. (2004). Hormonal regulation of plant growth and development. *PLoS Biology*, 2(9), e311.
- Gupta, K., Wani, S. H., Razzaq, A., Skalicky, M., Samantara, K., Gupta, S., Pandita, D., Goel, S., Grewal, S., Hejnak, V., Shiv, A., El-Sabrou, A. M., Elansary, H. O., Alaklabi, A. & Brestic, M. (2022). Absciscic Acid: Role in Fruit Development and Ripening. *Frontiers in Plant Science*, 13.
- Hedden, P. & Thomas, S. G. (2012). Gibberellin biosynthesis and its regulation. *Biochemical Journal*, 444(1), 11-25.
- Kieber, J. J. & Schaller, G. E. (2014). Cytokinins. *The Arabidopsis Book*, 12, e0168.
- Lupepsa, B. F., Ferrarese-Filho, O., Constantin, R. P., Marchiosi, R. & dos Santos, W. D. (2024). Gibberellin biosynthesis, its roles in plant physiology and abiotic stress responses. *Brazilian Journal of Development*.
- Nagel, J. C. (2024). Vegetais e hormônios.
- Peleg, Z. & Blumwald, E. (2011). Hormone balance and abiotic stress tolerance in crop plants. *Current Opinion in Plant Biology*, 14(3), 290-295.
- Rathod, S. A., Rabari, D. S. & Kansara, S. S. (2025). Plant Growth Promoting Rhizobacteria: Mechanisms and their Role in Sustainable Agriculture. *Biological Forum – An International Journal*, 17(1), 90-94.
- Singh, A., Singh, M. & Singh, P. (2023). Influence of Plant Growth Regulators on Root, Shoot and Yield Attributes of Strawberry (*Fragaria x ananassa* Duch.) under Protected Conditions. *Biological Forum – An International Journal*, 15(11), 136-142.
- Santner, A. & Estelle, M. (2009). Recent advances and emerging trends in plant hormone signaling. *Nature*, 459(7245), 1071-1078.

- Sasaki, A., Ashikari, M., Ueguchi-Tanaka, M., Itoh, H., Nishimura, A., Swapan, D. & Matsuoka, M. (2002). Green revolution: A mutant gibberellin-synthesis gene in rice. *Nature*, 416(6882), 701-702.
- Thapa, U., Ansari, Z. G., Ramesh, S., Krishnaveni, A. & Rabi, A. (2024). Plant Hormones and Growth Regulators: Mechanisms. *Interactions, and Agricultural Applications*, 3(3), 11–20.
- Zhu, J. K. (2016). Abiotic stress signaling and responses in plants. *Cell*, 167(2), 313-324.
- Zhao, J., Wang, J., Liu, B., Zhang, P., Kudoyarova, G. R., Liu, C. & Zhang, K. (2024). Spatially Distributed Cytokinins: Metabolism, Signaling, and Transport. *Plant Communications*, 100936.

How to cite this article: Achanta Rama (2025). From Seed to Harvest: How Plant Hormones Drive Growth and Agricultural Innovation. *AgriBio Innovations*, 2(1): 44-48.